



Contents lists available at ScienceDirect

Journal of Mathematical Analysis and Applications

www.elsevier.com/locate/jmaa


When does the k -hyponormality of a 2-variable weighted shift become subnormality? ☆

Jasang Yoon

Department of Mathematics, The University of Texas-Pan American, Edinburg, TX 78539, United States

ARTICLE INFO

Article history:

Received 15 November 2010

Available online 18 January 2011

Submitted by R. Curto

Keywords:

Jointly hyponormal pairs

Subnormal pairs

k -Hyponormal

2-variable weighted shift

Tensor core

ABSTRACT

In this article we construct a sequence of nontrivial classes of 2-variable weighted shifts $\{\mathcal{G}_k\}_{k=2}^{\infty}$ such that the k -hyponormality of an arbitrary power of a member $W_{(\alpha, \beta)}$ from \mathcal{G}_k is equivalent to its subnormality.

© 2011 Elsevier Inc. All rights reserved.

1. Introduction

Let \mathcal{H} be a complex Hilbert space and let $\mathcal{B}(\mathcal{H})$ denote the algebra of bounded linear operators on \mathcal{H} . For $S, T \in \mathcal{B}(\mathcal{H})$ let $[S, T] := ST - TS$. We say that an n -tuple $\mathbf{T} = (T_1, \dots, T_n)$ of operators on \mathcal{H} is (jointly) *hyponormal* if the operator matrix

$$[\mathbf{T}^*, \mathbf{T}] := \begin{pmatrix} [T_1^*, T_1] & [T_2^*, T_1] & \cdots & [T_n^*, T_1] \\ [T_1^*, T_2] & [T_2^*, T_2] & \cdots & [T_n^*, T_2] \\ \vdots & \vdots & \ddots & \vdots \\ [T_1^*, T_n] & [T_2^*, T_n] & \cdots & [T_n^*, T_n] \end{pmatrix}$$

is positive on the direct sum of n copies of \mathcal{H} (cf. [1,10]). The n -tuple \mathbf{T} is said to be *normal* if \mathbf{T} is commuting and each T_i is normal, and \mathbf{T} is *subnormal* if \mathbf{T} is the restriction of a normal n -tuple to a common invariant subspace. For $k \geq 1$, a commuting pair $\mathbf{T} \equiv (T_1, T_2)$ is said to be *k -hyponormal* if

$$\mathbf{T}(k) := (T_1, T_2, T_1^2, T_2 T_1, T_2^2, \dots, T_1^k, T_2 T_1^{k-1}, \dots, T_2^k)$$

is hyponormal, or equivalently

$$[\mathbf{T}(k)^*, \mathbf{T}(k)] = ([T_2^q T_1^p]^*, T_2^m T_1^n)_{\substack{1 \leq n+m \leq k \\ 1 \leq p+q \leq k}} \geq 0.$$

Clearly, normal \Rightarrow subnormal $\Rightarrow k$ -hyponormal. The Bram–Halmos criterion states that an operator $T \in \mathcal{B}(\mathcal{H})$ is subnormal if and only if the k -tuple (T, T^2, \dots, T^k) is hyponormal for all $k \geq 1$.

☆ This work was partially supported by a Faculty Research Council Grant at The University of Texas-Pan American.

E-mail address: yoonyj@utpa.edu.

URL: <http://www.math.utpa.edu/~yoonyj/>.

For $\alpha \equiv \{\alpha_n\}_{n=0}^\infty$ a bounded sequence of positive real numbers (called *weights*), let $W_\alpha : \ell^2(\mathbb{Z}_+) \rightarrow \ell^2(\mathbb{Z}_+)$ be the associated unilateral weighted shift, defined by $W_\alpha e_n := \alpha_n e_{n+1}$ (all $n \geq 0$), where $\{e_n\}_{n=0}^\infty$ is the canonical orthonormal basis in $\ell^2(\mathbb{Z}_+)$. We let $W_\alpha \equiv \text{shift}(\alpha_0, \alpha_1, \dots)$ and $U_+ := \text{shift}(1, 1, \dots)$ (the (unweighted) unilateral shift). For $0 < a \leq 1$ we also let $S_a := \text{shift}(a, 1, 1, \dots)$. The moments of W_α are given as

$$\gamma_k \equiv \gamma_k(W_\alpha) := \begin{cases} 1, & \text{if } k = 0 \\ \alpha_0^2 \cdots \alpha_{k-1}^2, & \text{if } k > 0 \end{cases}.$$

It is easy to see that W_α is never normal, and that it is hyponormal if and only if $\alpha_0 \leq \alpha_1 \leq \dots$. Similarly, consider double-indexed positive bounded sequences $\alpha_{\mathbf{k}}, \beta_{\mathbf{k}} \in \ell^\infty(\mathbb{Z}_+^2)$, $\mathbf{k} \equiv (k_1, k_2) \in \mathbb{Z}_+^2 := \mathbb{Z}_+ \times \mathbb{Z}_+$ and let $\ell^2(\mathbb{Z}_+^2)$ be the Hilbert space of square-summable complex sequences indexed by \mathbb{Z}_+^2 . (Recall that $\ell^2(\mathbb{Z}_+^2)$ is canonically isometrically isomorphic to $\ell^2(\mathbb{Z}_+) \otimes \ell^2(\mathbb{Z}_+)$.) We define the 2-variable weighted shift $W_{(\alpha, \beta)} \equiv (T_1, T_2)$ by

$$T_1 e_{\mathbf{k}} := \alpha_{\mathbf{k}} e_{\mathbf{k} + \varepsilon_1},$$

$$T_2 e_{\mathbf{k}} := \beta_{\mathbf{k}} e_{\mathbf{k} + \varepsilon_2},$$

where $\varepsilon_1 := (1, 0)$ and $\varepsilon_2 := (0, 1)$. Clearly,

$$T_1 T_2 = T_2 T_1 \Leftrightarrow \beta_{\mathbf{k} + \varepsilon_1} \alpha_{\mathbf{k}} = \alpha_{\mathbf{k} + \varepsilon_2} \beta_{\mathbf{k}} \quad (\text{all } \mathbf{k} \in \mathbb{Z}_+^2). \quad (1.1)$$

In an entirely similar way one can define multivariable weighted shifts. Trivially, a pair of unilateral weighted shifts W_α and W_β gives rise to a 2-variable weighted shift $W_{(\alpha, \beta)}$, if we let $\alpha_{(k_1, k_2)} := \alpha_{k_1}$ and $\beta_{(k_1, k_2)} := \beta_{k_2}$ (all $k_1, k_2 \in \mathbb{Z}_+$). In this case, $W_{(\alpha, \beta)}$ is subnormal (resp. hyponormal) if and only if so are T_1 and T_2 ; in fact, under the canonical identification of $\ell^2(\mathbb{Z}_+^2)$ with $\ell^2(\mathbb{Z}_+) \otimes \ell^2(\mathbb{Z}_+)$, we have $T_1 \cong I \otimes W_\alpha$ and $T_2 \cong W_\beta \otimes I$, and $W_{(\alpha, \beta)}$ is also doubly commuting. For this reason, we do not focus attention on shifts of this type, and use them only when the above mentioned triviality is desirable or needed. Given $\mathbf{k} \equiv (k_1, k_2) \in \mathbb{Z}_+^2$, the moments of $W_{(\alpha, \beta)}$ of order \mathbf{k} are

$$\gamma_{\mathbf{k}} \equiv \gamma_{\mathbf{k}}(W_{(\alpha, \beta)}) := \begin{cases} 1, & \text{if } (k_1, k_2) = (0, 0), \\ \alpha_{(0,0)}^2 \cdots \alpha_{(k_1-1,0)}^2, & \text{if } k_1 \geq 1 \text{ and } k_2 = 0, \\ \beta_{(0,0)}^2 \cdots \beta_{(0,k_2-1)}^2, & \text{if } k_1 = 0 \text{ and } k_2 \geq 1, \\ \alpha_{(0,0)}^2 \cdots \alpha_{(k_1-1,0)}^2 \beta_{(k_1,0)}^2 \cdots \beta_{(k_1,k_2-1)}^2, & \text{if } k_1 \geq 1 \text{ and } k_2 \geq 1. \end{cases} \quad (1.2)$$

We remark that, due to the commutativity condition (1.1), $\gamma_{\mathbf{k}}$ can be computed using any nondecreasing path from $(0, 0)$ to (k_1, k_2) . We now recall a well-known characterization of subnormality for multivariable weighted shifts [20], due to C. Berger (cf. [2, III.8.16]) and independently established by R. Gellar and L.J. Wallen [18] in the single variable case: $W_{(\alpha, \beta)}$ admits a commuting normal extension if and only if there is a probability measure μ (which we call the *Berger measure* of $W_{(\alpha, \beta)}$) defined on the 2-dimensional rectangle $R = [0, a_1] \times [0, a_2]$ (where $a_i := \|T_i\|^2$) such that $\gamma_{\mathbf{k}} = \int_R \mathbf{t}^{\mathbf{k}} d\mu(\mathbf{t}) := \int_R t_1^{k_1} t_2^{k_2} d\mu(\mathbf{t})$, for all $\mathbf{k} \in \mathbb{Z}_+^2$. Observe that U_+ and S_a are subnormal, with Berger measures δ_1 and $(1 - a^2)\delta_0 + a^2\delta_1$, respectively, where δ_p denotes the point-mass probability measure with support the singleton set $\{p\}$. Single and multivariable weighted shifts have played an important role in the study of the problems of existence of commuting normal extensions (cf. [5–7, 9, 13–15, 21, 26]). They have also played a significant role in the study of cyclicity and reflexivity, in the study of C^* -algebras generated by multiplication operators on Bergman spaces, as fertile ground to test new hypotheses, and as canonical models for theories of dilation and positivity (cf. [12, 19, 22]). We need some further notation to describe our results. We use \mathfrak{H}_0 (resp. \mathfrak{H}_∞) to denote the set of commuting pairs of subnormal operators (resp. subnormal pairs) on Hilbert space. For $k \geq 1$, we let \mathfrak{H}_k denote the class of k -hyponormal pairs in \mathfrak{H}_0 . Clearly, $\mathfrak{H}_\infty \subseteq \cdots \subseteq \mathfrak{H}_k \subseteq \cdots \subseteq \mathfrak{H}_2 \subseteq \mathfrak{H}_1 \subseteq \mathfrak{H}_0$. The main results in [5, 13] show that these inclusions are all proper. For an arbitrary 2-variable weighted shift $W_{(\alpha, \beta)}$, we let \mathcal{M}_i (resp. \mathcal{N}_j) be the subspace of $\ell^2(\mathbb{Z}_+^2)$ which is spanned by the canonical orthonormal basis associated to indices $\mathbf{k} \equiv (k_1, k_2)$ with $k_1 \geq 0$ and $k_2 \geq i$ (resp. $k_1 \geq j$ and $k_2 \geq 0$). We will often write \mathcal{M}_1 simply as \mathcal{M} and \mathcal{N}_1 as \mathcal{N} . The core $c(W_{(\alpha, \beta)})$ of $W_{(\alpha, \beta)}$ is the restriction of $W_{(\alpha, \beta)}$ to the invariant subspace $\mathcal{M} \cap \mathcal{N}$. A 2-variable weighted shift $W_{(\alpha, \beta)}$ is said to be of *tensor form* if it is of the form $(I \otimes W_\alpha, W_\beta \otimes I)$. The class of all 2-variable weighted shifts $W_{(\alpha, \beta)} \in \mathfrak{H}_0$ whose core is of tensor form will be denoted by \mathcal{TC} ; in symbols, $\mathcal{TC} := \{W_{(\alpha, \beta)} \in \mathfrak{H}_0 : c(W_{(\alpha, \beta)}) \text{ is of tensor form}\}$. Note that if $W_{(\alpha, \beta)} \in \mathcal{TC}$, then $W_{(\alpha, \beta)} \in \mathfrak{H}_\infty$. Given a subnormal 2-variable weighted shift $W_{(\alpha, \beta)}$ with Berger measure μ , we let $W_{\alpha^{(j)}}$ ($j \geq 0$) (resp. $W_{\beta^{(i)}}$ ($i \geq 0$)) denote the associated j -th horizontal (resp. i -th vertical) slice of $W_{(\alpha, \beta)}$. Clearly, $W_{\alpha^{(j)}}$ (resp. $W_{\beta^{(i)}}$) is subnormal, and we let ξ_j (resp. η_i) denote its Berger measure. For $k \geq 1$ we let $\mathcal{G}_k := \{W_{(\alpha, \beta)} \in \mathcal{A} : W_{(\alpha, \beta)}|_{\mathcal{M}} \equiv (I \otimes S_a, U_+ \otimes I) \text{ and } \text{card}(\text{supp } \xi_0) \leq (k+1)\}$, where $\mathcal{A} := \{W_{(\alpha, \beta)} \in \mathcal{TC} : c(W_{(\alpha, \beta)}) \text{ has 1-atomic Berger measure}\}$ and $\text{card}(\text{supp } \xi_0)$ means the cardinality of the support of ξ_0 (cf. see Fig. 1(ii)). For the meaning of set inclusion, we clearly have $\mathcal{G}_1 = \mathcal{S}_1 \subsetneq \mathcal{G}_2 \subsetneq \cdots \subsetneq \mathcal{G}_k \subsetneq \cdots \subsetneq \mathcal{A} \subsetneq \mathcal{TC}$. Observe that a 2-variable weighted shift $W_{(\alpha, \beta)} \in \mathcal{S}_1$ has a core with Berger measure $\delta_{\{1,1\}} = \delta_1 \times \delta_1$. For $k \geq 1$, we note that if $W_{(\alpha, \beta)} \in \mathcal{G}_k$, then $W_{(\alpha, \beta)}$ can be fully determined by 3 parameters: the weight $a := \alpha_{(0,1)}$, the weight $y := \beta_{(0,0)}$ and the Berger measure ξ_0 of the 0-th horizontal subnormal slice $\text{shift}(x_0, x_1, x_2, \dots)$ of $W_{(\alpha, \beta)}$. Thus we can denote a 2-variable weighted shift $W_{(\alpha, \beta)} \in \mathcal{G}_k$ by $\langle a, y, \xi_0 \rangle$.




Fig. 1. Weight diagram of a generic 2-variable weighted shift in S_1 and the 2-variable weighted shift in Theorem 2.5, respectively.

2. Main results

For a general operator T on Hilbert space, it is well known that the subnormality of T implies the subnormality of T^h ($h \geq 2$). The converse implication, however, is false; in fact, the subnormality of all powers T^h ($h \geq 2$) does not necessarily imply the subnormality of T , even if $T \equiv W_\alpha$ is a unilateral weighted shift [24, p. 378]. Thus, it is natural to ask when the subnormality of all powers W_α^h ($h \geq 2$) does imply the subnormality of W_α . More general, we might ask when the k -hyponormality ($k \geq 1$) of all powers W_α^h ($h \geq 2$) does imply the subnormality of W_α . For $k \geq 2$, we let $W_\alpha \equiv \text{shift}(a, b, 1, 1, \dots)$ where $0 < a < b < 1$. Then W_α^h ($h \geq 2$) is subnormal, but W_α is not k -hyponormal ($k \geq 2$), because the k -hyponormality of W_α ($k \geq 2$) $\Leftrightarrow b = 1$. For $k = 1$, we consider $W_\alpha \equiv \text{shift}(1, 1 - x, y, y, \dots)$ where $0 < x < 1 < y$. Then a simple calculation shows that W_α^h ($h \geq 2$) is subnormal, but W_α is not hyponormal. In the multivariable case, we can consider these analogous results. The standard assumption on a pair $\mathbf{T} \equiv (T_1, T_2)$ is that each component T_i is subnormal ($i = 1, 2$). With this in mind, the analogous questions are highly nontrivial. In [8,17], we identified a large nontrivial class S_1 (cf. see Fig. 1(i)) of 2-variable weighted shifts for which the 2-hyponormality of an arbitrary power of the initial pair is equivalent to subnormality of the initial pair. Thus, it is natural to consider

Problem 2.1. (See [17, Problem 6.8].) Is S_1 the largest class in \mathcal{A} for which the implication

$$W_{(\alpha, \beta)}^{(h_0, \ell_0)} := (T_1^{h_0}, T_2^{\ell_0}) \in \bigoplus \mathfrak{H}_2 \quad \text{for some } h_0, \ell_0 \geq 1 \quad \Rightarrow \quad W_{(\alpha, \beta)} \in \mathfrak{H}_\infty$$

holds?

In this paper we give a concrete answer for Problem 2.1 above and build a class \mathcal{G}_k ($k \geq 2$) in \mathcal{A} such that if $W_{(\alpha, \beta)} \in \mathcal{G}_k$ with $\text{card}(\text{supp } \xi_0) = k + 1$, then for some $h_0, \ell_0 \geq 1$

$$W_{(\alpha, \beta)}^{(h_0, \ell_0)} \in \bigoplus \mathfrak{H}_k \quad \Leftrightarrow \quad W_{(\alpha, \beta)} \in \mathfrak{H}_\infty.$$

For this, we first recall that, in one variable, the n -th power of a weighted shift is unitarily equivalent to the direct sum of n weighted shifts. Something similar happens in two variables, as we will see in the proof of Theorem 2.5 below. We let $\mathcal{H} \equiv \ell^2(\mathbb{Z}_+) = \bigvee_{j=0}^\infty \{e_j\}$. Given integers i and h ($h \geq 1$, $0 \leq i \leq h-1$), define $\mathcal{H}_i := \bigvee_{j=0}^\infty \{e_{hj+i}\}$; clearly, $\mathcal{H} = \bigoplus_{i=0}^{h-1} \mathcal{H}_i$. Following the notation in [11], for a weight sequence $\alpha \equiv \{\alpha_n\}_{n=0}^\infty$ we let

$$W_{\alpha(h:i)} := \text{shift} \left(\prod_{n=0}^{h-1} \alpha_{hj+i+n} \right)_{j=0}^\infty; \quad (2.1)$$

that is, $W_{\alpha(h:i)}$ denotes the sequence of products of weights in adjacent packets of size h , beginning with $\alpha_i \cdots \alpha_{i+h-1}$. For example, given a weighted shift $W_\alpha \equiv \text{shift}(\alpha_0, \alpha_1, \dots)$, we have $W_{\alpha(2:0)} = \text{shift}(\alpha_0\alpha_1, \alpha_2\alpha_3, \dots)$, $W_{\alpha(2:1)} = \text{shift}(\alpha_1\alpha_2, \alpha_3\alpha_4, \dots)$ and $W_{\alpha(3:2)} = \text{shift}(\alpha_2\alpha_3\alpha_4, \alpha_5\alpha_6\alpha_7, \dots)$. For $h \geq 1$, $0 \leq i \leq h-1$, we note that $W_{\alpha(h:i)}$ is unitarily equivalent to $W_\alpha|_{\mathcal{H}_i}$. Therefore, W_α^h is unitarily equivalent to $\bigoplus_{i=0}^{h-1} W_{\alpha(h:i)}$. Consequently, if W_α is subnormal with the

Berger measure μ , then $W_{\alpha(h,i)}$ is subnormal with the Berger measure μ_i , where

$$d\mu_0(s) = d\mu(s^{\frac{1}{h}}) \quad \text{and} \quad d\mu_i(s) = \frac{s^{\frac{1}{h}}}{\gamma_i} d\mu(s^{\frac{1}{h}}) \quad \text{for } 1 \leq i \leq h-1. \quad (2.2)$$

Furthermore, we have

Lemma 2.2. (See [11, Corollary 2.8].)

- (i) Let $k \geq 1$. Then W_{α}^h is k -hyponormal $\Leftrightarrow W_{\alpha(h,i)}$ is k -hyponormal for $0 \leq i \leq h-1$.
- (ii) W_{α}^h is subnormal $\Leftrightarrow W_{\alpha(h,i)}$ is subnormal for $0 \leq i \leq h-1$.

We now introduce a key family of examples for our main results. For $k \geq 2$, $0 < a_i$ ($1 \leq i \leq k-1$), $0 \leq c_0, c_1, \dots, c_k \leq 1$ (with $\sum_{i=0}^k c_i = 1$), we let $W_x \equiv \text{shift}(x_0, x_1, \dots)$ be given by

$$x_n := \begin{cases} \sqrt{\sum_{i=1}^{k-1} c_i a_i + c_k}, & \text{if } n = 0, \\ \sqrt{\frac{\sum_{i=1}^{k-1} c_i a_i^{n+1} + c_k}{\sum_{i=1}^{k-1} c_i a_i^n + c_k}}, & \text{if } n \geq 1. \end{cases} \quad (2.3)$$

We now consider

$$\xi_0 := c_0 \delta_0 + \sum_{i=1}^{k-1} c_i \delta_{a_i} + c_k \delta_1. \quad (2.4)$$

Then ξ_0 is a probability measure. For $n = 0$, we denote $\gamma_0(W_x)$ by 1. We note that for $n \geq 1$ the moments associated with W_x are

$$\begin{aligned} \gamma_n(W_x) &\equiv x_0^2 x_1^2 x_2^2 \cdots x_{n-1}^2 \\ &= \left(\sum_{i=1}^{k-1} c_i a_i + c_k \right) \cdot \left(\frac{\sum_{i=1}^{k-1} c_i a_i^2 + c_k}{\sum_{i=1}^{k-1} c_i a_i + c_k} \right) \cdots \left(\frac{\sum_{i=1}^{k-1} c_i a_i^n + c_k}{\sum_{i=1}^{k-1} c_i a_i^{n-1} + c_k} \right) \\ &= \sum_{i=1}^{k-1} c_i a_i^n + c_k = \int s^n d\xi_0(s) \quad (n \geq 1). \end{aligned}$$

Thus, it follows that W_x is subnormal, with Berger measure

$$d\xi_0(s) = c_0 d\delta_0(s) + \sum_{i=1}^{k-1} c_i d\delta_{a_i}(s) + c_k d\delta_1(s).$$

Lemma 2.3. For $k \geq 2$, $0 < a_i$ ($1 \leq i \leq k-1$), $0 \leq c_0, c_1, \dots, c_k \leq 1$ (with $\sum_{i=0}^k c_i = 1$), $h_0 \geq 1$, $k_1 \geq 0$, we let

$$G(h_0, k_1, k) := \begin{pmatrix} \sum_{i=1}^{k-1} c_i a_i^{k_1+2h_0} + c_k & \sum_{i=1}^{k-1} c_i a_i^{k_1+3h_0} + c_k & \cdots & \sum_{i=1}^{k-1} c_i a_i^{k_1+(k+1)h_0} + c_k \\ \sum_{i=1}^{k-1} c_i a_i^{k_1+3h_0} + c_k & \sum_{i=1}^{k-1} c_i a_i^{k_1+4h_0} + c_k & \cdots & \sum_{i=1}^{k-1} c_i a_i^{k_1+(k+2)h_0} + c_k \\ \vdots & \vdots & \ddots & \vdots \\ \sum_{i=1}^{k-1} c_i a_i^{k_1+(k+1)h_0} + c_k & \sum_{i=1}^{k-1} c_i a_i^{k_1+(k+2)h_0} + c_k & \cdots & \sum_{i=1}^{k-1} c_i a_i^{k_1+2kh_0} + c_k \end{pmatrix}.$$

Then $G(h_0, k_1, k)$ is invertible and

$$\det G(h_0, k_1, k) = c_k \prod_{i=1}^{k-1} c_i a_i^{k_1+2h_0} (1 - a_i^{h_0})^2 \cdot \prod_{i < j}^{k-1} (a_i^{h_0} - a_j^{h_0})^2, \quad (2.5)$$

where $\prod_{i < j}^{k-1} (a_i^{h_0} - a_j^{h_0})^2 := 1$, if $k = 2$.

Proof. By a direct calculation using *Mathematica* [25], we have (2.5). The invertibility of $G(h_0, k_1, k)$ is clear from (2.5). \square

Lemma 2.4. Under the conditions of Lemma 2.3 and for $0 < a, y \leq 1$, we let

$$D(a, y) := \begin{pmatrix} y^2 & a^2 y^2 & y^2 \\ a^2 y^2 & a^2 y^2 & a^2 y^2 \\ y^2 & a^2 y^2 & 1 \end{pmatrix},$$

$$F(a, y, h_0, 0, k) := \begin{pmatrix} a^2 y^2 & a^2 y^2 & \cdots & a^2 y^2 \\ a^2 y^2 & a^2 y^2 & \cdots & a^2 y^2 \\ \sum_{i=1}^{k-1} c_i a_i^{h_0} + c_k & \sum_{i=1}^{k-1} c_i a_i^{2h_0} + c_k & \cdots & \sum_{i=1}^{k-1} c_i a_i^{kh_0} + c_k \end{pmatrix},$$

$$P(a, y, h_0, 0, k) := \begin{pmatrix} D(a, y) & F(a, y, h_0, 0, k) \\ F^*(a, y, h_0, 0, k) & G(h_0, 0, k) \end{pmatrix}.$$

Then we have

$$P(a, y, h_0, 0, k) \geq 0 \Leftrightarrow y \leq \begin{cases} \min\{\sqrt{\frac{c_k}{a^2}}, \sqrt{\frac{c_0}{1-a^2}}\}, & \text{if } 0 < a < 1, \\ \sqrt{c_k}, & \text{if } a = 1. \end{cases} \quad (2.6)$$

Proof. By Lemma 2.3, since $D(h_0)$ is invertible, if we apply Lemma A.5 to $P(a, y, h_0, 0, k)$, we have

$$P(a, y, h_0, 0, k) \geq 0 \Leftrightarrow D(a, y) - W(a, y, h_0, 0, k)^* G(h_0, 0, k) W(a, y, h_0, 0, k) \geq 0,$$

where

$$W(a, y, h_0, 0, k) := \begin{pmatrix} \frac{\prod_{i=1}^{k-1} \frac{a^2 y^2 a_i^{h_0}}{(a_i^{h_0} - 1) c_k}}{a^2 y^2 (a_1^{h_0} a_2^{h_0} \cdots a_{k-2}^{h_0} + \cdots + a_2^{h_0} a_3^{h_0} \cdots a_{k-1}^{h_0})} & \frac{\prod_{i=1}^{k-1} \frac{a^2 y^2 a_i^{h_0}}{(a_i^{h_0} - 1) c_k}}{a^2 y^2 (a_1^{h_0} a_2^{h_0} \cdots a_{k-2}^{h_0} + \cdots + a_2^{h_0} a_3^{h_0} \cdots a_{k-1}^{h_0})} & g_1(h_0, k) \\ \frac{(-1)^{\prod_{i=1}^{k-1} (a_i^{h_0} - 1) c_k}}{a^2 y^2 (a_1^{h_0} a_2^{h_0} \cdots a_{k-3}^{h_0} + \cdots + a_3^{h_0} a_4^{h_0} \cdots a_{k-1}^{h_0})} & \frac{(-1)^{\prod_{i=1}^{k-1} (a_i^{h_0} - 1) c_k}}{a^2 y^2 (a_1^{h_0} a_2^{h_0} \cdots a_{k-3}^{h_0} + \cdots + a_3^{h_0} a_4^{h_0} \cdots a_{k-1}^{h_0})} & g_2(h_0, k) \\ \frac{(-1)^2 \prod_{i=1}^{k-1} (a_i^{h_0} - 1) c_k}{(-1)^2 \prod_{i=1}^{k-1} (a_i^{h_0} - 1) c_k} & \frac{(-1)^2 \prod_{i=1}^{k-1} (a_i^{h_0} - 1) c_k}{(-1)^2 \prod_{i=1}^{k-1} (a_i^{h_0} - 1) c_k} & g_3(h_0, k) \\ \vdots & \vdots & \vdots \\ \frac{(-1)^{k-1} \frac{a^2 y^2 (\sum_{i=1}^{k-1} a_i^{h_0})}{\prod_{i=1}^{k-1} (a_i^{h_0} - 1) c_k}}{(-1)^k a^2 y^2} & \frac{(-1)^{k-1} \frac{a^2 y^2 (\sum_{i=1}^{k-1} a_i^{h_0})}{\prod_{i=1}^{k-1} (a_i^{h_0} - 1) c_k}}{(-1)^k a^2 y^2} & g_{k-1}(h_0, k) \\ \frac{(-1)^{k-1} \frac{a^2 y^2 (\sum_{i=1}^{k-1} a_i^{h_0})}{\prod_{i=1}^{k-1} (a_i^{h_0} - 1) c_k}}{(-1)^k a^2 y^2} & \frac{(-1)^{k-1} \frac{a^2 y^2 (\sum_{i=1}^{k-1} a_i^{h_0})}{\prod_{i=1}^{k-1} (a_i^{h_0} - 1) c_k}}{(-1)^k a^2 y^2} & g_k(h_0, k) \end{pmatrix}$$

and

$$g_1(h_0, k) := 1 + \sum_{i=1}^{k-1} a_i^{-h_0}, \quad g_2(h_0, k) := (-1) \left(\sum_{i=1}^{k-1} a_i^{-h_0} + \sum_{i < j}^{k-1} a_i^{-h_0} a_j^{-h_0} \right),$$

$$g_3(h_0, k) := (-1)^2 \left(\sum_{i < j}^{k-1} a_i^{-h_0} a_j^{-h_0} + \sum_{i < j < \ell}^{k-1} a_i^{-h_0} a_j^{-h_0} a_\ell^{-h_0} \right),$$

$$g_{k-1}(h_0, k) := (-1)^{k-1} \left(a_1^{-h_0} a_2^{-h_0} \cdots a_{k-2}^{-h_0} + \cdots + a_2^{-h_0} a_3^{-h_0} \cdots a_{k-1}^{-h_0} + \prod_{i=1}^{k-1} a_i^{-h_0} \right),$$

$$g_k(h_0, k) := (-1)^k \prod_{i=1}^{k-1} a_i^{-h_0}.$$

A direct calculation shows that

$$D(a, y) - W(a, y, h_0, 0, k)^* G(h_0, 0, k) W(a, y, h_0, 0, k) \geq 0$$

$$\Leftrightarrow \begin{pmatrix} y^2 - \frac{a^4 y^4}{c_k} & a^2 y^2 - \frac{a^4 y^4}{c_k} & (1 - a^2) y^2 \\ a^2 y^2 - \frac{a^4 y^4}{c_k} & a^2 y^2 - \frac{a^4 y^4}{c_k} & 0 \\ (1 - a^2) y^2 & 0 & c_0 \end{pmatrix} \geq 0$$

$$\Leftrightarrow \begin{pmatrix} y^2 - \frac{a^4 y^4}{c_k} + \frac{(1-a^2)^2 y^4}{c_0} & a^2 y^2 - \frac{a^4 y^4}{c_k} \\ a^2 y^2 - \frac{a^4 y^4}{c_k} & a^2 y^2 - \frac{a^4 y^4}{c_k} \end{pmatrix} =: \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} =: A \geq 0.$$

To check $A \geq 0$, it is sufficient to check $a_{22} \geq 0$ and $\det A \geq 0$, because $(a_{22} \geq 0 \text{ and } \det A \geq 0) \Rightarrow a_{11} \geq 0$. Thus, a straightforward calculation shows that $a_{22} \geq 0 \Leftrightarrow a^2 y^2 \leq c_k$ and

$$\det A \geq 0 \Leftrightarrow (-y^2 + a^2 y^2 + c_0)(c_k - a^2 y^2) \geq 0.$$

Thus, it follows that

$$\begin{aligned} P(a, y, h_0, 0, k) \geq 0 &\Leftrightarrow \{a^2 y^2 \leq c_{k-1} \text{ and } y^2(1 - a^2) \leq c_0\} \\ &\Leftrightarrow \begin{cases} \min\{\sqrt{\frac{c_k}{a^2}}, \sqrt{\frac{c_0}{1-a^2}}\}, & \text{if } 0 < a < 1, \\ \sqrt{c_k}, & \text{if } a = 1. \end{cases} \quad \square \end{aligned}$$

For our main results, we recall that for $k \geq 1$, $\mathcal{G}_k = \{W_{(\alpha, \beta)} \in \mathcal{A} : W_{(\alpha, \beta)}|_{\mathcal{M}} \equiv (I \otimes S_a, U_+ \otimes I) \text{ and } \text{card}(\text{supp } \xi_0) \leq (k+1)\}$ (cf. see Fig. 1(ii)). We then have

Theorem 2.5. For $k \geq 2$, we let $W_{(\alpha, \beta)} \equiv (T_1, T_2) \equiv (a, y, \xi_0) \in \mathcal{G}_k$ (where the 0-th horizontal slice $W_x \equiv \text{shift}(x_0, x_1, \dots)$ is as in (2.3) with $0 < c_0, c_1, \dots, c_k < 1$). Then the following statements are equivalent:

- (i) $W_{(\alpha, \beta)} \in \mathfrak{H}_k$;
- (ii) $W_{(\alpha, \beta)} \in \mathfrak{H}_\infty$;
- (iii) for some $h_0, \ell_0 \geq 1$, $W_{(\alpha, \beta)}^{(h_0, \ell_0)} \equiv (T_1^{h_0}, T_2^{\ell_0}) \in \bigoplus \mathfrak{H}_k$.

Proof. (i) \Rightarrow (ii): From Lemma A.2, we recall that a 2-variable weighted shift $W_{(\alpha, \beta)}$ is k -hyponormal if and only if

$$M_{\mathbf{k}}(k) = (\gamma_{\mathbf{k}+(m,n)+(p,q)})_{\substack{0 \leq n+m \leq k \\ 0 \leq p+q \leq k}} \geq 0,$$

for all $\mathbf{k} \in \mathbb{Z}_+^2$. It is straightforward to verify that $W_{(\alpha, \beta)}|_{\mathcal{M}} \cong (I \otimes S_a, U_+ \otimes I)$, so that $W_{(\alpha, \beta)}|_{\mathcal{M}}$ is subnormal. Since $W_{(\alpha, \beta)} \in \mathfrak{H}_k$, if we apply Lemma A.2(ii) at $\mathbf{k} = (k_1, 0)$ (all $k_1 \geq 0$), we have $M_{(k_1, 0)}(k) \geq 0$. We note that for $h_0, \ell_0 \geq 1$ the moments $\gamma_{\mathbf{k}}(\mathbf{k} \in \mathbb{Z}_+^2)$ associated with $W_{(\alpha, \beta)}^{(h_0, \ell_0)}$ are

$$\gamma_{\mathbf{k}}(W_{(\alpha, \beta)}^{(h_0, \ell_0)}) = \begin{cases} 1, & \text{if } k_1 = 0 \text{ and } k_2 = 0, \\ \gamma_{k_1 h_0}(W_\alpha), & \text{if } k_1 \geq 1 \text{ and } k_2 = 0, \\ y^2, & \text{if } k_1 = 0 \text{ and } k_2 \geq 1, \\ a^2 y^2, & \text{if } k_1 \geq 1 \text{ and } k_2 \geq 1. \end{cases} \quad (2.7)$$

Thus, by (2.7), we have that

$$W_{(\alpha, \beta)}^{(1, 1)} \equiv W_{(\alpha, \beta)} \in \mathfrak{H}_k \Rightarrow M_{(k_1, 0)}(k)(W_{(\alpha, \beta)}) \geq 0 \Rightarrow M_{(0, 0)}(k)(W_{(\alpha, \beta)}) \geq 0.$$

From Lemma 2.4, a direct computation (i.e., interchanging rows and columns, discarding a redundant row and column in the moment matrix of $M_{(0, 0)}(k)(W_{(\alpha, \beta)})$) shows that

$$\begin{aligned} M_{(0, 0)}(k)(W_{(\alpha, \beta)}) \geq 0 &\Leftrightarrow P(a, y, 1, 0, k) \geq 0 \\ &\Leftrightarrow y \leq \begin{cases} \min\{\sqrt{\frac{c_k}{a^2}}, \sqrt{\frac{c_0}{1-a^2}}\}, & \text{if } 0 < a < 1, \\ \sqrt{c_k}, & \text{if } a = 1. \end{cases} \end{aligned}$$

Thus, we have

$$W_{(\alpha, \beta)} \in \mathfrak{H}_k \Rightarrow y \leq \begin{cases} \min\{\sqrt{\frac{c_k}{a^2}}, \sqrt{\frac{c_0}{1-a^2}}\}, & \text{if } 0 < a < 1, \\ \sqrt{c_k}, & \text{if } a = 1. \end{cases} \quad (2.8)$$

We now characterize the subnormality of $W_{(\alpha, \beta)}$ using its parametric characterizations. Lemma A.3 will help us characterize $W_{(\alpha, \beta)} \in \mathfrak{H}_\infty$. Since $W_{(\alpha, \beta)}|_{\mathcal{M}} \equiv (I \otimes S_a, U_+ \otimes I)$ is subnormal with Berger measure $\mu_{\mathcal{M}} \equiv [(1 - a^2)\delta_0 + a^2\delta_1] \times \delta_1$, we can think of $W_{(\alpha, \beta)}$ as a backward extension of $W_{(\alpha, \beta)}|_{\mathcal{M}}$ (in the t direction) and apply Lemma A.3. Note that $d(\mu_{\mathcal{M}})_{\text{ext}}(s, t) \equiv [(1 - a^2)\delta_0 + a^2\delta_1] \times \delta_1$ and $\alpha_{00}^2 \|\frac{1}{t}\|_{L^1(\mu_{\mathcal{M}})}(\mu_{\mathcal{M}})_{\text{ext}}^X = y^2[(1 - a^2)\delta_0 + a^2\delta_1]$. From Lemma 2.4 observe that

$$P(a, y, 1, 0, k) \geq 0 \Leftrightarrow y^2[(1-a^2)\delta_0 + a^2\delta_1] \leq c_0\delta_0 + \sum_{i=1}^{k-1} c_i\delta_{a_i} + c_k\delta_1$$

$$\Leftrightarrow \alpha_{00}^2 \left\| \frac{1}{t} \right\|_{L^1(\mu_{\mathcal{M}})} (\mu_{\mathcal{M}})_{\text{ext}}^X \leq \xi_0, \quad \text{where } \xi_0 \text{ is the Berger measure of } W_x \text{ as in (2.3).}$$

Thus, by Lemma A.3, we have

$$W_{(\alpha, \beta)} \in \mathfrak{H}_\infty \Leftrightarrow P(a, y, 1, 0, k) \geq 0.$$

Therefore, by Lemma 2.4 and (2.8), it follows that $W_{(\alpha, \beta)} \in \mathfrak{H}_k \Rightarrow W_{(\alpha, \beta)} \in \mathfrak{H}_\infty$.

(ii) \Rightarrow (iii): Since $W_{(\alpha, \beta)} \in \mathfrak{H}_\infty$, for all $h_0, \ell_0 \geq 1$, we have $W_{(\alpha, \beta)}^{(h_0, \ell_0)} \in \bigoplus \mathfrak{H}_\infty$ by the functional calculus. Thus we get for some $h_0, \ell_0 \geq 1$, $W_{(\alpha, \beta)}^{(h_0, \ell_0)} \in \bigoplus \mathfrak{H}_k$.

(iii) \Rightarrow (i): For some $h_0, \ell_0 \geq 1$, we suppose that $W_{(\alpha, \beta)}^{(h_0, \ell_0)} \in \bigoplus \mathfrak{H}_k$. Fixed $h_0, \ell_0 \geq 1$, we first let $\mathcal{H}_{(m, n)} := \bigvee_{i, j=0}^\infty \{e_{(h_0 i + m, \ell_0 j + n)} : h_0, \ell_0 \geq 1\}$, for $0 \leq m \leq h_0 - 1$ and $0 \leq n \leq \ell_0 - 1$. Then we have

$$\ell^2(\mathbb{Z}_+^2) \equiv \bigoplus_{m=0}^{h_0-1} \bigoplus_{n=0}^{\ell_0-1} \mathcal{H}_{(m, n)}.$$

Observe that $\mathcal{H}_{(m, n)}$ reduces $T_1^{h_0}$ and $T_2^{\ell_0}$. Thus, if a 2-variable weighted shift $W_{(\alpha, \beta)}$ is given in Fig. 1(ii), then for $h_0, \ell_0 \geq 1$, we can write

$$W_{(\alpha, \beta)}^{(h_0, \ell_0)} \equiv (T_1^{h_0}, T_2^{\ell_0})$$

$$\cong (W_{\alpha(h_0:0)} \oplus (I \otimes S_a), T_2|_{\mathcal{H}_0}) \oplus \bigoplus_{i=1}^{h_0-1} (W_{\alpha(h_0:i)} \oplus (I \otimes U_+), T_2|_{\mathcal{H}_i}) \quad [\text{cf. see Fig. 2(i)}],$$

where

$$W_{\alpha(h_0:i)} = \text{shift} \left(\sqrt{\frac{\gamma_{(i+1)h_0}}{\gamma_{ih_0}}}, \sqrt{\frac{\gamma_{(i+2)h_0}}{\gamma_{(i+1)h_0}}}, \dots \right) \quad \text{and} \quad \mathcal{H}_i := \bigoplus_{n=0}^{\ell_0-1} \mathcal{H}_{(i, n)} \quad (0 \leq i \leq k-1).$$

Thus, we observe that $W_{(\alpha, \beta)}^{(h_0, \ell_0)} \in \bigoplus \mathfrak{H}_k$ is equivalent to $(W_{\alpha(h_0:0)} \oplus (I \otimes S_a), T_2|_{\mathcal{H}_0}) \in \mathfrak{H}_k$ and $(W_{\alpha(h_0:i)} \oplus (I \otimes U_+), T_2|_{\mathcal{H}_i}) \in \mathfrak{H}_k$, for $1 \leq i \leq h_0 - 1$. To show $W_{(\alpha, \beta)} \in \mathfrak{H}_k$, by Lemma A.2, it is enough to show that $W_{(\alpha, \beta)}|_{\mathcal{M}} \in \mathfrak{H}_k$, $W_{(\alpha, \beta)}|_{\mathcal{N}} \in \mathfrak{H}_k$ and $M_{(0,0)}(k)(W_{(\alpha, \beta)}) \geq 0$. Since $W_{(\alpha, \beta)}|_{\mathcal{M}} \cong (I \otimes S_a, U_+ \otimes I)$ is subnormal, we need to show $W_{(\alpha, \beta)}|_{\mathcal{N}} \in \mathfrak{H}_k$ and $M_{(0,0)}(k)(W_{(\alpha, \beta)}) \geq 0$. For $W_{(\alpha, \beta)}|_{\mathcal{N}} \in \mathfrak{H}_k$, we first want to show that for some $h_0, \ell_0 \geq 1$

$$(W_{(\alpha, \beta)}|_{\mathcal{N}})^{(h_0, \ell_0)} \in \bigoplus \mathfrak{H}_k \Leftrightarrow W_{(\alpha, \beta)}|_{\mathcal{N}} \in \mathfrak{H}_k. \quad (2.9)$$

Since $W_{(\alpha, \beta)}|_{\mathcal{M} \cap \mathcal{N}} \cong (I \otimes U_+, U_+ \otimes I)$ is subnormal, to show (2.9), we need to show that

$$M_{(k_1, 0)}(k)(W_{(\alpha, \beta)}^{(h_0, \ell_0)}) \geq 0 \Leftrightarrow M_{(k_1, 0)}(k)(W_{(\alpha, \beta)}) \geq 0 \quad \text{for } k_1 \geq 1.$$

Since $G(h_0, k_1, k)$ is invertible, by (2.7), Fig. 2(ii) and Lemma A.5, for $k_1 \geq 1$ and some $h_0, \ell_0 \geq 1$, we have that

$$M_{(k_1, 0)}(k)(W_{(\alpha, \beta)}^{(h_0, \ell_0)}) \geq 0 \Leftrightarrow M \geq 0 \Leftrightarrow a^2 y^2 \leq c_k \Leftrightarrow M_{(k_1, 0)}(2)(W_{(\alpha, \beta)}) \geq 0,$$

where

$$M := \begin{pmatrix} \begin{pmatrix} a^2 y^2 & a^2 y^2 \\ a^2 y^2 & \sum_{i=1}^{k-1} c_i a_i^{h_0} + c_k \end{pmatrix} & \begin{pmatrix} a^2 y^2 & \dots & a^2 y^2 \\ \sum_{i=1}^{k-1} c_i a_i^{k_1+h_0} + c_k & \dots & \sum_{i=1}^{k-1} c_i a_i^{k_1+kh_0} + c_k \end{pmatrix} \\ \begin{pmatrix} a^2 y^2 & \sum_{i=1}^{k-1} c_i a_i^{k_1+h_0} + c_k \\ \vdots & \vdots \\ a^2 y^2 & \sum_{i=1}^{k-1} c_i a_i^{k_1+kh_0} + c_k \end{pmatrix} & G(h_0, k_1, k) \end{pmatrix}.$$

Thus, we have

$$(W_{(\alpha, \beta)}|_{\mathcal{N}})^{(h_0, \ell_0)} \in \bigoplus \mathfrak{H}_k \Leftrightarrow W_{(\alpha, \beta)}|_{\mathcal{N}} \in \mathfrak{H}_k.$$




Fig. 2. Weight diagram of the 2-variable weighted shift $W_{(\alpha, \beta)}^{(h_0, \ell_0)}$ in Theorem 2.5 and weight diagram of the 2-variable weighted shift in Lemma A.4, respectively.

For $M_{(0,0)}(k)(W_{(\alpha, \beta)}) \geq 0$, we note that

$$W_{(\alpha, \beta)}^{(h_0, \ell_0)} \in \bigoplus \mathfrak{H}_k \Rightarrow (W_{\alpha(h_0:0)} \oplus (I \otimes S_a), T_2|_{\mathcal{H}_0}) \in \mathfrak{H}_k.$$

We also observe that

$$(W_{\alpha(h_0:0)} \oplus (I \otimes S_a), T_2|_{\mathcal{H}_0}) \cong \bigoplus_{n=0}^{\ell_0-1} (W_{\alpha(h_0:0)} \oplus (I \otimes S_a), T_2^{\ell_0}|_{\mathcal{H}_{(0,n)}})$$

and

$$\bigoplus_{n=0}^{\ell_0-1} (W_{\alpha(h_0:0)} \oplus (I \otimes S_a), T_2^{\ell_0}|_{\mathcal{H}_{(0,n)}}) \cong (W_{\alpha(h_0:0)} \oplus (I \otimes S_a), T_2^{\ell_0}|_{\mathcal{H}_{(0,0)}}) \oplus \bigoplus_{n=0}^{\ell_0-1} (I \otimes S_a, U_+ \otimes I).$$

Note that the second summand is clearly subnormal; thus, for $h_0, \ell_0 \geq 1$, the k -hyponormality of $(W_{\alpha(h_0:0)} \oplus (I \otimes S_a), T_2|_{\mathcal{H}_0})$ is equivalent to the k -hyponormality of the first summand, $(W_{\alpha(h_0:0)} \oplus (I \otimes S_a), T_2^{\ell_0}|_{\mathcal{H}_{(0,0)}})$. Observe also that

$$(W_{\alpha(h_0:0)} \oplus (I \otimes S_a), T_2^{\ell_0}|_{\mathcal{H}_{(0,0)}}) \cong (W_{\alpha(h_0:0)} \oplus (I \otimes S_a), T_2|_{\mathcal{H}_{(0,0)}}).$$

Thus we have

$$(W_{\alpha(h_0:0)} \oplus (I \otimes S_a), T_2|_{\mathcal{H}_0}) \in \mathfrak{H}_k \Leftrightarrow (W_{\alpha(h_0:0)} \oplus (I \otimes S_a), T_2|_{\mathcal{H}_{(0,0)}}) \in \mathfrak{H}_k.$$

To check the k -hyponormality of $(W_{\alpha(h_0:0)} \oplus (I \otimes S_a), T_2|_{\mathcal{H}_{(0,0)}})$, we observe that it suffices to apply Lemma A.2(ii) at $\mathbf{k} = (0, 0)$. From Lemma 2.4 and (2.7), after we apply Lemma A.2 to $(0, 0)$ of $(W_{\alpha(h_0:0)} \oplus (I \otimes U_+), T_2|_{\mathcal{H}_0})$, we have

$$M_{(0,0)}(k)(W_{\alpha(h_0:0)} \oplus (I \otimes U_+), T_2|_{\mathcal{H}_0}) \geq 0 \Leftrightarrow P(a, y, h_0, 0, k) \geq 0 \Leftrightarrow y^2 \leq c_2,$$

where $P(a, y, h_0, 0, k)$ is as in Lemma 2.4. Thus we get

$$W_{(\alpha, \beta)}^{(h_0, \ell_0)} \in \bigoplus \mathfrak{H}_k \Rightarrow y^2 \leq c_k \Leftrightarrow M_{(0,0)}(k)(W_{(\alpha, \beta)}) \geq 0. \quad (2.10)$$

Therefore, we have for some $h_0, \ell_0 \geq 1$,

$$W_{(\alpha, \beta)}^{(h_0, \ell_0)} \in \bigoplus \mathfrak{H}_k \Rightarrow W_{(\alpha, \beta)} \in \mathfrak{H}_k$$

and our proof is now complete. \square

Remark 2.6.

- (i) We note that $\text{card}(\text{supp } \xi_0) = (k+1)$, where ξ_0 is as in Theorem 2.5.
 (ii) In Theorem 2.5, we show that for given $k \geq 2$, there exists a 2-variable weighted shift $W_{(\alpha, \beta)} \in \mathcal{G}_k \subsetneq \mathcal{A}$ for which some $h_0, \ell_0 \geq 1$

$$W_{(\alpha, \beta)}^{(h_0, \ell_0)} \in \bigoplus \mathfrak{H}_k \Leftrightarrow W_{(\alpha, \beta)} \in \mathfrak{H}_\infty.$$

- (iii) By Theorem 2.5, we note that there exists a 2-variable weighted shift $W_{(\alpha, \beta)} \in \mathcal{G}_2$ in \mathcal{A} for which some $h_0, \ell_0 \geq 1$

$$W_{(\alpha, \beta)}^{(h_0, \ell_0)} \in \bigoplus \mathfrak{H}_k \Leftrightarrow W_{(\alpha, \beta)} \in \mathfrak{H}_\infty.$$

Thus, Theorem 2.5 gives an answer for Problem 2.1 and more.

Observe that if $W_{(\alpha, \beta)} \equiv \langle a, y, \xi_0 \rangle \in \mathcal{G}_k$ ($k \geq 2$), then by (2.4) the measure ξ_0 can be completely determined by the $(2k-1)$ parameters $\{a_i\}_{i=1}^{k-1}$ and $\{c_i\}_{i=0}^k$. Thus we can also denote a 2-variable weighted shift $W_{(\alpha, \beta)} \equiv \langle a, y, \xi_0 \rangle \in \mathcal{G}_k$ by $\langle a, y, \{a_i\}_{i=1}^{k-1}, \{c_i\}_{i=0}^k \rangle$ (cf. see Fig. 1(ii)). We now assume $ay \leq \sum_{i=1}^{k-1} c_i a_i^h + c_k$ (all $h \geq 1$), because we need to ensure that $W_{(\alpha, \beta)} \in \mathfrak{H}_0$. We now obtain a canonical representation for the powers $\langle a, y, \{a_i\}_{i=1}^{k-1}, \{c_i\}_{i=0}^k \rangle^{(h, \ell)}$ as an orthogonal direct sum of 2-variable weighted shifts in \mathcal{G}_k . In what follows, we abbreviate the orthogonal direct sums of h copies of a shift $\langle a, y, \{a_i\}_{i=1}^{k-1}, \{c_i\}_{i=0}^k \rangle$ by $h \cdot \langle a, y, \{a_i\}_{i=1}^{k-1}, \{c_i\}_{i=0}^k \rangle$. Then we have

Proposition 2.7. We let $\langle a, y, \{a_i\}_{i=1}^{k-1}, \{c_i\}_{i=0}^k \rangle \equiv \langle a, y, \{a_i\}_{i=1}^{k-1}, c_0, \{c_i\}_{i=1}^{k-1}, c_k \rangle \in \mathcal{G}_k$. Then for $h, \ell \geq 1$, we have

$$\begin{aligned} & \langle a, y, \{a_i\}_{i=1}^{k-1}, c_0, \{c_i\}_{i=1}^{k-1}, c_k \rangle^{(h, \ell)} \\ & \cong \langle a, y, \{a_i^{\frac{1}{h}}\}_{i=1}^{k-1}, c_0, \{c_i\}_{i=1}^{k-1}, c_k \rangle \\ & \oplus \left\langle 1, \frac{ay}{\sqrt{\sum_{i=1}^{k-1} c_i a_i^{mh} + c_k}}, \{a_i^{\frac{1}{h}}\}_{i=1}^{k-1}, 0, \left\{ \frac{a_i^{\frac{1}{h}} c_i}{\sqrt{\sum_{i=1}^{k-1} c_i a_i^m + c_k}} \right\}_{i=1}^{k-1}, \frac{c_k}{\sqrt{\sum_{i=1}^{k-1} c_i a_i^m + c_k}} \right\rangle \\ & \oplus (\ell-1) \cdot \langle a, 1, 0, 1-a^2, 0, a^2 \rangle \oplus (h-1)(\ell-1) \cdot \langle 1, 1, 0, 0, 0, 1 \rangle, \end{aligned}$$

so that the class \mathcal{G}_k is invariant under all powers.

Proof. We recall that we decompose the space $\ell^2(\mathbb{Z}_+^2)$ as the orthogonal direct sum of $h\ell$ subspaces $\mathcal{H}_{(m,n)}$, each isometrically isomorphic to $\ell^2(\mathbb{Z}_+^2)$, namely $\mathcal{H}_{(m,n)} := \bigvee_{i,j=0}^\infty \{e_{(hi+m, \ell j+n)}\}$ ($0 \leq m \leq h-1$, $0 \leq n \leq \ell-1$). This particular decomposition allows us to write the power

$$\langle a, y, \{a_i\}_{i=1}^{k-1}, c_0, \{c_i\}_{i=1}^{k-1}, c_k \rangle^{(h, \ell)}$$

as the orthogonal direct sum

$$\bigoplus_{0 \leq m \leq h-1, 0 \leq n \leq \ell-1}^{(h, \ell)} \langle a, y, \{a_i\}_{i=1}^{k-1}, c_0, \{c_i\}_{i=1}^{k-1}, c_k \rangle|_{\mathcal{H}_{(m,n)}}.$$

From (2.2), we will now identify each of the summands $\langle a, y, b, c_0, c_1, c_2 \rangle^{(h, \ell)}|_{\mathcal{H}_{(m,n)}}$ ($0 \leq m \leq h-1$, $0 \leq n \leq \ell-1$).

Case 1: ($m=0, n=0$) Direct inspection of the weight families α and β shows that

$$\langle a, y, \{a_i\}_{i=1}^{k-1}, c_0, \{c_i\}_{i=1}^{k-1}, c_k \rangle^{(h, \ell)} e_{(hi, \ell j)} = \langle a, y, \{a_i^{\frac{1}{h}}\}_{i=1}^{k-1}, c_0, \{c_i\}_{i=1}^{k-1}, c_k \rangle e_{(hi, \ell j)},$$

and therefore

$$\langle a, y, \{a_i\}_{i=1}^{k-1}, c_0, \{c_i\}_{i=1}^{k-1}, c_k \rangle^{(h, \ell)}|_{\mathcal{H}_{(0,0)}} \cong \langle a, y, \{a_i^{\frac{1}{h}}\}_{i=1}^{k-1}, c_0, \{c_i\}_{i=1}^{k-1}, c_k \rangle.$$

Case 2: ($m > 0, n = 0$) In this case the generic basis vector of $\mathcal{H}_{(m,0)}$ is $e_{(hi+m,\ell j)}$, so that

$$\begin{aligned} & \langle a, y, \{a_i\}_{i=1}^{k-1}, c_0, \{c_i\}_{i=1}^{k-1}, c_k \rangle^{(h,\ell)} e_{(hi+m,\ell j)} \\ &= \left\langle 1, \frac{ay}{\sqrt{\sum_{i=1}^{k-1} c_i a_i^{mh} + c_k}}, \{a_i^{\frac{1}{h}}\}_{i=1}^{k-1}, 0, \left\{ \frac{a_i^{\frac{1}{h}} c_i}{\sqrt{\sum_{i=1}^{k-1} c_i a_i^m + c_k}} \right\}_{i=1}^{k-1}, \frac{c_k}{\sqrt{\sum_{i=1}^{k-1} c_i a_i^m + c_k}} \right\rangle e_{(hi+m,\ell j)}. \end{aligned}$$

It follows that

$$\begin{aligned} & \langle a, y, \{a_i\}_{i=1}^{k-1}, c_0, \{c_i\}_{i=1}^{k-1}, c_k \rangle^{(h,\ell)} |_{\mathcal{H}_{(m,0)}} \\ & \cong \left\langle 1, \frac{ay}{\sqrt{\sum_{i=1}^{k-1} c_i a_i^{mh} + c_k}}, \{a_i^{\frac{1}{h}}\}_{i=1}^{k-1}, 0, \left\{ \frac{a_i^{\frac{1}{h}} c_i}{\sqrt{\sum_{i=1}^{k-1} c_i a_i^m + c_k}} \right\}_{i=1}^{k-1}, \frac{c_k}{\sqrt{\sum_{i=1}^{k-1} c_i a_i^m + c_k}} \right\rangle. \end{aligned}$$

Case 3: ($m = 0, n > 0$) In this case the generic basis vector of $\mathcal{H}_{(0,n)}$ is $e_{(hi,\ell j+n)}$, and therefore

$$\langle a, y, \{a_i\}_{i=1}^{k-1}, c_0, \{c_i\}_{i=1}^{k-1}, c_k \rangle^{(h,\ell)} e_{(hi,\ell j+n)} = \langle a, 1, 0, 1 - a^2, 0, a^2 \rangle e_{(hi,\ell j+n)}.$$

It follows that $\langle a, y, \{a_i\}_{i=1}^{k-1}, c_0, \{c_i\}_{i=1}^{k-1}, c_k \rangle^{(h,\ell)} |_{\mathcal{H}_{(0,n)}} \cong \langle a, 1, 0, 1 - a^2, 0, a^2 \rangle$.

Case 4: ($m > 0, n > 0$) Since $\mathcal{H}_{(m,n)} \subseteq \mathcal{M} \cap \mathcal{N}$, and the core of $W_{(\alpha,\beta)}$ is trivial, it is clear that all relevant weights are equal to 1, so

$$\langle a, y, \{a_i\}_{i=1}^{k-1}, c_0, \{c_i\}_{i=1}^{k-1}, c_k \rangle^{(h,\ell)} e_{(hi+m,\ell j+n)} = \langle 1, 1, 0, 0, 0, 1 \rangle e_{(hi+m,\ell j+n)},$$

and therefore $\langle a, y, \{a_i\}_{i=1}^{k-1}, c_0, \{c_i\}_{i=1}^{k-1}, c_k \rangle^{(h,\ell)} |_{\mathcal{H}_{(m,n)}} \cong \langle 1, 1, 0, 0, 0, 1 \rangle$.

Therefore, our proof is now complete. \square

Corollary 2.8. For $\langle a, y, \{a_i\}_{i=1}^{k-1}, \{c_i\}_{i=0}^k \rangle \in \mathcal{G}_k$ ($k \geq 2$), the following statements are equivalent:

- (i) for some $h_0, \ell_0 \geq 1$, $\langle a, y, \{a_i\}_{i=1}^{k-1}, \{c_i\}_{i=0}^k \rangle^{(h_0,\ell_0)} \in \bigoplus \mathfrak{H}_k$;
- (ii) for all $h, \ell \geq 1$, $\langle a, y, \{a_i\}_{i=1}^{k-1}, \{c_i\}_{i=0}^k \rangle^{(h,\ell)} \in \bigoplus \mathfrak{H}_k$;
- (iii) $\langle a, y, \{a_i\}_{i=1}^{k-1}, \{c_i\}_{i=0}^k \rangle \in \mathfrak{H}_k$;
- (iv) for some $h_0, \ell_0 \geq 1$, $\langle a, y, \{a_i\}_{i=1}^{k-1}, \{c_i\}_{i=0}^k \rangle^{(h_0,\ell_0)} \in \bigoplus \mathfrak{H}_\infty$;
- (v) for all $h, \ell \geq 1$, $\langle a, y, \{a_i\}_{i=1}^{k-1}, \{c_i\}_{i=0}^k \rangle^{(h,\ell)} \in \bigoplus \mathfrak{H}_\infty$;
- (vi) $\langle a, y, \{a_i\}_{i=1}^{k-1}, \{c_i\}_{i=0}^k \rangle \in \mathfrak{H}_\infty$.

Proof. This is straightforward from Theorem 2.5, Proposition 2.7 and the functional calculus. \square

Acknowledgments

The author is deeply indebted to the referee for a number of comments and observations that helped improve the content and presentation of this paper. The portions of the proof of some results were obtained using calculations with the software tool *Mathematica* [25].

Appendix A

For the reader's convenience, in this section we gather several well-known auxiliary results which are needed for the proofs of the main results in this article. First, to detect hyponormality for 2-variable weighted shifts we use a simple criterion involving a base point \mathbf{k} in \mathbb{Z}_+^2 and its five neighboring points in $\mathbf{k} + \mathbb{Z}_+^2$ at path distance at most 2.

Lemma A.1 (Six-point Test). (See [3, Theorem 6.1].) Let $W_{(\alpha,\beta)} \equiv (T_1, T_2)$ be a 2-variable weighted shift, with weight sequences α and β . Then

$$\begin{aligned} & [W_{(\alpha,\beta)}^*, W_{(\alpha,\beta)}] \geq 0 \\ \Leftrightarrow & H(k_1, k_2)(1) := \begin{pmatrix} \alpha_{\mathbf{k}+\varepsilon_1}^2 - \alpha_{\mathbf{k}}^2 & \alpha_{\mathbf{k}+\varepsilon_2} \beta_{\mathbf{k}+\varepsilon_1} - \alpha_{\mathbf{k}} \beta_{\mathbf{k}} \\ \alpha_{\mathbf{k}+\varepsilon_2} \beta_{\mathbf{k}+\varepsilon_1} - \alpha_{\mathbf{k}} \beta_{\mathbf{k}} & \beta_{\mathbf{k}+\varepsilon_2}^2 - \beta_{\mathbf{k}}^2 \end{pmatrix} \geq 0 \quad (\text{for all } \mathbf{k} \in \mathbb{Z}_+^2). \end{aligned}$$

Next, we present an analogous criterion to detect the k -hyponormality of 2-variable weighted shifts.

Lemma A.2. (See [5, Theorem 2.4].) Let $W_{(\alpha,\beta)}$ be a 2-variable weighted shift with weight sequence α and β . The following statements are equivalent:

- (i) $W_{(\alpha,\beta)}$ is k -hyponormal;
- (ii) $M_{\mathbf{k}}(\mathbf{k}) := (\gamma_{\mathbf{k}+(n,m)+(p,q)})_{\substack{0 \leq n+m \leq k \\ 0 \leq p+q \leq k}} \geq 0$ for all $\mathbf{k} \in \mathbb{Z}_+^2$.

To check subnormality of 2-variable weighted shifts, we introduce some definitions.

- (i) Let μ and ν be two positive measures on \mathbb{R}_+ . We say that $\mu \leq \nu$ on $X := \mathbb{R}_+$, if $\mu(E) \leq \nu(E)$ for all Borel subset $E \subseteq \mathbb{R}_+$; equivalently, $\mu \leq \nu$ if and only if $\int f d\mu \leq \int f d\nu$ for all $f \in C(X)$ such that $f \geq 0$ on \mathbb{R}_+ .
- (ii) Let μ be a probability measure on $X \times Y$, and assume that $\frac{1}{t} \in L^1(\mu)$. The extremal measure μ_{ext} (which is also a probability measure) on $X \times Y$ is given by $d\mu_{\text{ext}}(s, t) := (1 - \delta_0(t)) \frac{1}{t \|\frac{1}{t}\|_{L^1(\mu)}} d\mu(s, t)$.
- (iii) Given a measure μ on $X \times Y$, the marginal measure μ^X is given by $\mu^X := \mu \circ \pi_X^{-1}$, where $\pi_X : X \times Y \rightarrow X$ is the canonical projection onto X . Thus $\mu^X(E) = \mu(E \times Y)$, for every $E \subseteq X$.

Then we have:

Lemma A.3 (Subnormal backward extension). (See [13, Proposition 3.10].) Let $W_{(\alpha,\beta)}$ be a 2-variable weighted shift, and assume that $W_{(\alpha,\beta)}|_{\mathcal{M}}$ is subnormal with associated measure $\mu_{\mathcal{M}}$ and that $W_0 := \text{shift}(\alpha_0, \alpha_{10}, \dots)$ is subnormal with associated measure ξ_0 . Then $W_{(\alpha,\beta)}$ is subnormal if and only if

- (i) $\frac{1}{t} \in L^1(\mu_{\mathcal{M}})$;
- (ii) $\beta_{00}^2 \leq (\|\frac{1}{t}\|_{L^1(\mu_{\mathcal{M}})})^{-1}$;
- (iii) $\beta_{00}^2 \|\frac{1}{t}\|_{L^1(\mu_{\mathcal{M}})} (\mu_{\mathcal{M}})_{\text{ext}}^X \leq \xi_0$.

Moreover, if $\beta_{00}^2 \|\frac{1}{t}\|_{L^1(\mu_{\mathcal{M}})} = 1$, then $(\mu_{\mathcal{M}})_{\text{ext}}^X = \xi_0$. In the case when $W_{(\alpha,\beta)}$ is subnormal, the Berger measure μ of $W_{(\alpha,\beta)}$ is given by

$$d\mu(s, t) = \beta_{00}^2 \left\| \frac{1}{t} \right\|_{L^1(\mu_{\mathcal{M}})} d(\mu_{\mathcal{M}})_{\text{ext}}(s, t) + \left(d\xi_0(s) - \beta_{00}^2 \left\| \frac{1}{t} \right\|_{L^1(\mu_{\mathcal{M}})} d(\mu_{\mathcal{M}})_{\text{ext}}^X(s) \right) d\delta_0(t).$$

Lemma A.4. (See [27, Theorem 2.8].) Let $W_{(\alpha,\beta)} \in \mathfrak{H}_0$ be a 2-variable weighted shift whose weight diagram is given in Fig. 2(ii), so that $W_{(\alpha,\beta)}|_{\mathcal{M}} \cong (I \otimes \text{shift}(\beta_1, \beta_2, \dots), U_+ \otimes bI)$. Assume that $\|W_\alpha\| = b > 0$, where $W_\alpha \equiv \text{shift}(\alpha_0, \alpha_1, \alpha_2, \dots)$. Then $W_{(\alpha,\beta)} \in \mathfrak{H}_1 \Leftrightarrow W_{(\alpha,\beta)} \in \mathfrak{H}_\infty \Leftrightarrow$ the Berger measure μ_α of W_α has an atom at b^2 .

Lemma A.5. (Cf. [4, Proposition 2.2], [16,23].) Let $M \equiv \begin{pmatrix} A & B \\ B^* & C \end{pmatrix}$ be a 2×2 operator matrix, where A and C are square matrices and B is a rectangular matrix. Then

$$M \geq 0 \Leftrightarrow \text{there exists } W \text{ such that } \begin{cases} A \geq 0, \\ B = AW, \\ C \geq W^*AW. \end{cases}$$

References

- [1] A. Athavale, On joint hyponormality of operators, Proc. Amer. Math. Soc. 103 (1988) 417–423.
- [2] J. Conway, The Theory of Subnormal Operators, Math. Surveys Monogr., vol. 36, Amer. Math. Soc., Providence, 1991.
- [3] R. Curto, Joint hyponormality: A bridge between hyponormality and subnormality, Proc. Sympos. Pure Math. 51 (1990) 69–91.
- [4] R. Curto, L. Fialkow, Solution of the Truncated Complex Moment Problem with Flat Data, Mem. Amer. Math. Soc., vol. 568, Amer. Math. Soc., Providence, 1996.
- [5] R. Curto, S.H. Lee, J. Yoon, k -Hyponormality of multivariable weighted shifts, J. Funct. Anal. 229 (2005) 462–480.
- [6] R. Curto, S.H. Lee, J. Yoon, Hyponormality and subnormality for powers of commuting pairs of subnormal operators, J. Funct. Anal. 245 (2007) 390–412.
- [7] R. Curto, S.H. Lee, J. Yoon, Reconstruction of the Berger measure when the core is of tensor form, in: Actas del XVI Coloquio Latinoamericano de Álgebra, Bibl. Rev. Mat. Iberoamericana (2007) 317–331.
- [8] R. Curto, S.H. Lee, J. Yoon, Which 2-hyponormal 2-variable weighted shifts are subnormal?, Linear Algebra Appl. 429 (2008) 2227–2238.
- [9] R. Curto, S.H. Lee, J. Yoon, New approach to the 2-variable subnormal completion problem, J. Math. Anal. Appl. 370 (2010) 270–283.
- [10] R. Curto, P. Muhly, J. Xia, Hyponormal pairs of commuting operators, Oper. Theory Adv. Appl. 35 (1988) 1–22.
- [11] R. Curto, S. Park, k -Hyponormality of powers of weighted shifts, Proc. Amer. Math. Soc. 131 (2002) 2762–2769.
- [12] R. Curto, M. Putinar, Nearly subnormal operators and moments problems, J. Funct. Anal. 115 (1993) 480–497.
- [13] R. Curto, J. Yoon, Jointly hyponormal pairs of subnormal operators need not be jointly subnormal, Trans. Amer. Math. Soc. 358 (2006) 5139–5159.
- [14] R. Curto, J. Yoon, Disintegration-of-measure techniques for multivariable weighted shifts, Proc. London Math. Soc. 93 (2006) 381–402.

- [15] R. Curto, J. Yoon, Propagation phenomena for hyponormal 2-variable weighted shifts, *J. Operator Theory* 58 (1) (2007) 101–130.
- [16] R. Curto, J. Yoon, Spectral picture of 2-variable weighted shifts, *C. R. Acad. Sci. Paris* 343 (2006) 579–584.
- [17] R. Curto, J. Yoon, When is hyponormality for 2-variable weighted shifts invariant under powers?, *Indiana Univ. Math. J.*, submitted for publication.
- [18] R. Gellar, L.J. Wallen, Subnormal weighted shifts and the Halmos–Bram criterion, *Proc. Japan Acad.* 46 (1970) 375–378.
- [19] P.R. Halmos, *A Hilbert Space Problem Book*, second ed., *Grad. Texts in Math.*, vol. 19, Springer-Verlag, New York/Berlin, 1982.
- [20] N.P. Jewell, A.R. Lubin, Commuting weighted shifts and analytic function theory in several variables, *J. Operator Theory* 1 (1979) 207–223.
- [21] A.R. Lubin, Weighted shifts and commuting normal extension, *J. Austral. Math. Soc. Ser. A* 27 (1) (1979) 17–26.
- [22] A.L. Shields, Weighted shift operators and analytic function theory, *Math. Surveys* 13 (1974) 49–128.
- [23] Ju.L. Smul'jan, An operator Hellinger integral, *Mat. Sb. (N.S.)* 49 (1959) 381–430 (in Russian).
- [24] J. Stampfli, Which weighted shifts are subnormal?, *Pacific J. Math.* 17 (1966) 367–379.
- [25] Wolfram Research, Inc., *Mathematica*, Version 4.2, Wolfram Research, Inc., Champaign, IL, 2002.
- [26] J. Yoon, Disintegration of measures and contractive 2-variable weighted shifts, *Integral Equations Operator Theory* 59 (2007) 281–298.
- [27] J. Yoon, Schur product techniques for commuting multivariable weighted shifts, *J. Math. Anal. Appl.* 333 (2007) 626–641.